


# Frequency-domain ray series for viscoelastic waves with a non-symmetric stiffness matrix

*Luděk Klimeš*

Department of Geophysics  
Faculty of Mathematics and Physics  
Charles University in Prague



<http://sw3d.cz>

Symmetry of the frequency-domain elastic or viscoelastic stiffness tensor:  
 $c^{ijkl} = c^{ijkl}(x^m, \omega)$ :

$$c^{ijkl} = c^{jikl} \quad c^{ijkl} = c^{ijlk}$$

Additional symmetry of the frequency-domain stiffness tensor proved in an elastic medium but not in a viscoelastic medium:

$$c^{ijkl} = c^{klij}$$

Frequency-domain viscoelastic stiffness tensor:

$$c^{ijkl} \neq c^{klij}$$

We thus propose the frequency-domain ray series for viscoelastic waves with a stiffness tensor which is non-symmetric with respect to the exchange of the first pair of indices and the second pair of indices.

The lower-case Roman indices take values 1, 2 and 3. The Einstein summation over repetitive lower-case Roman indices is used hereinafter.

Frequency-domain viscoelastodynamic equation for complex-valued displacement  $u_i = u_i(x^m, \omega)$ :

$$(c^{ijkl} u_{l,k})_{,j} - (i\omega)^2 \rho u_i = 0 \quad (5)$$

Lower-case Roman subscript  $_{,k}$  following a comma denotes the partial derivative with respect to corresponding spatial coordinate  $x^k$ .

$\rho = \rho(x^m)$ ... density,

$\omega$ ... circular frequency.

Displacement in terms of its amplitude  $U_i = u_i(x^m, \omega)$  and travel time  $\tau = \tau(x^m)$ :

$$u_i = U_i \exp(i\omega\tau) \quad (6)$$

High-frequency asymptotic series

$$U_i = \sum_{n=0}^{\infty} (i\omega)^{-n} U_i^{[n]} \quad (7)$$

We consider standard anisotropic ray theory assuming strictly decoupled S waves, and proceed according to Červený (2001) using differential operators

$$N^i(U_m, \tau, n) = \varrho [\Gamma^{il}(x^m, \tau, n) U_l - U_i] \quad (9)$$

$$M^i(U_m, \tau, n) = (c^{ijkl} \tau_{,k} U_l)_{,j} + c^{ijkl} \tau_{,j} U_{l,k} \quad (10)$$

$$L^i(U_m) = (c^{ijkl} U_{l,k})_{,j} \quad (11)$$

Christoffel matrix

$$\Gamma^{il}(x^m, p_n) = c^{ijkl}(x^m) p_j p_k [\varrho(x^n)]^{-1} \quad (12)$$

is a function of six phase-space coordinates  $x^m, p_n$  formed by three spatial coordinates  $x^m$  and three slowness-vector components  $p_n$ .

The Christoffel matrix is not symmetric. Its right-hand eigenvectors differ from its left-hand eigenvectors.

Right-hand eigenvector  $g_i = g_i(x^m, \tau, n)$ , corresponding to selected eigenvalue  $G = G(x^m, \tau, n)$  of the Christoffel matrix:

$$\Gamma^{il} g_l = G g_i \quad (16)$$

Corresponding left-hand eigenvector  $\vec{g}_i = \vec{g}_i(x^m, \tau, n)$ :

$$\vec{g}_i \Gamma^{il} = \vec{g}_i G \quad (17)$$

We denote by  $G^\perp$  the other two eigenvalues of the Christoffel matrix, by  $g_i^\perp$  the corresponding right-hand eigenvectors, and by  $\vec{g}_i^\perp$  the corresponding left-hand eigenvectors. Superscript  $^\perp$  takes two values. The three right-hand eigenvectors of the Christoffel matrix and the three left-hand eigenvectors of the Christoffel matrix are mutually biorthogonal, and we choose them mutually biorthonormal.

Eikonal equation

$$G(x^m, \tau, n) = 1 \quad (20)$$

can be solved by the standard methods developed for solving the Hamilton-Jacobi equation (Hamilton, 1837; Červený, 1972; Klimeš, 2002; 2010).

Decomposition of a vectorial amplitude into principal amplitude component  $U_i^{[n]}$  and two additional amplitude components  $U^\perp[n]$ :

$$U_i^{[n]} = U^{[n]} g_i + \sum_{\perp} U^\perp[n] g_i^\perp \quad (30)$$

Additional amplitude components:

$$U^\perp[n] = -\varrho^{-1} \left[ \vec{g}_i^\perp M^i (U_k^{[n-1]}, \tau, n) + \vec{g}_i^\perp L^i (U_k^{[n-2]}) \right] (G^\perp - 1)^{-1} \quad (32)$$

with both  $U^{\perp[0]} = 0$ .

Zero-order principal amplitude component:

$$U^{[0]} = U_0^{[0]} (\varrho_0 J_0)^{\frac{1}{2}} (\varrho J)^{-\frac{1}{2}} \exp\left(\int_{\tau_0}^{\tau} d\gamma S\right) \quad (40)$$

Subscript  $_0$  denotes the initial conditions.

Squared geometrical spreading

$$J = \det\left(\frac{\partial x^i}{\partial \gamma^a}\right) \quad (41)$$

represents the Jacobian of transformation from ray coordinates  $\gamma^1, \gamma^2, \gamma^3$  to spatial coordinates  $x^i$ . These ray coordinates are composed of ray parameters  $\gamma^1$  and  $\gamma^2$ , and of travel time  $\gamma^3 = \tau$  along rays.

Difference between symmetric and non-symmetric stiffness matrices:

$$\begin{aligned}
 \mathbf{S} = & \frac{1}{4} \sum_{\perp} \left( \vec{g}_k \frac{\partial \Gamma^{kl}}{\partial x^j} g_l^{\perp} \vec{g}_r^{\perp} \frac{\partial \Gamma^{rs}}{\partial p_j} g_s - \vec{g}_k \frac{\partial \Gamma^{kl}}{\partial p_j} g_l^{\perp} \vec{g}_r^{\perp} \frac{\partial \Gamma^{rs}}{\partial x^j} g_s \right) (G - G^{\perp})^{-1} \\
 & - \frac{1}{4\rho} \vec{g}_i (c^{ijkl} - c^{ikjl})_{,j} \tau_{,k} g_l - \vec{g}_i \frac{dg_i}{d\gamma}
 \end{aligned} \tag{55}$$

Term  $\vec{g}_i \frac{dg_i}{d\gamma}$  represents just the correction of principal amplitude  $U^{[n]}$  due to the undefined length of right-hand eigenvector  $g_i$ , and may be put to zero.

Quantity  $S$  may be singular at slowness-surface singularities, but is regular at spatial caustics.

Quantity  $S$  vanishes for a symmetric stiffness matrix. For a non-symmetric stiffness matrix, quantity  $S$  vanishes in a homogeneous medium.

Higher-order principal amplitude components:

$$U^{[n]} = U^{[0]} \left[ \frac{U_0^{[n]}}{U_0^{[0]}} + \int_{\tau_0}^{\tau} d\gamma \frac{Z^{[n-1]}}{U^{[0]} \sqrt{\varrho}} \right] \quad (42)$$

with

$$Z^{[n-1]} = -\frac{1}{2\sqrt{\varrho}} \left[ \sum_{\perp} \vec{g}_i M^i (U^{\perp[n]} g_k^{\perp}, \tau, n) + \vec{g}_i L^i (U_k^{[n-1]}) \right] \quad (39)$$




## References:

- Červený, V. (1972): Seismic rays and ray intensities in inhomogeneous anisotropic media. *Geophys. J. R. astr. Soc.*, **29**, 1–13.
- Červený, V. (2001): *Seismic Ray Theory*. Cambridge Univ. Press, Cambridge.
- Hamilton, W.R. (1837): Third supplement to an essay on the theory of systems of rays. *Trans. Roy. Irish Acad.*, **17**, 1–144, read January 23, 1832, and October 22, 1832.
- Klimeš, L. (2002): Second-order and higher-order perturbations of travel time in isotropic and anisotropic media. *Stud. geophys. geod.*, **46**, 213–248.
- Klimeš, L. (2010): Transformation of spatial and perturbation derivatives of travel time at a general interface between two general media. *Seismic Waves in Complex 3-D Structures*, **20**, 103–114, online at “<http://sw3d.cz>”.
- Klimeš, L. (2016): Frequency-domain ray series for viscoelastic waves with a non-symmetric stiffness matrix. *Seismic Waves in Complex 3-D Structures*, **26**, 159–166, online at “<http://sw3d.cz>”.

# Acknowledgements

The research has been supported:

by the Grant Agency of the Czech Republic under contract 16-05237S,  
and by the consortium “Seismic Waves in Complex 3-D Structures”



<http://sw3d.cz>